Burden of disease profile

For the percentage of deaths by broad cause, deaths are classified into four groups namely: (i) injuries; (ii) non-communicable diseases; (iii) HIV and TB; and (iv) communicable diseases together with maternal perinatal and nutritional conditions. Data are given by gender and age group for the period 2010–2015. The second part of the graph shows the 10 leading single causes of death within each age group (both genders) for 2010–2015 combined.

Percentage of deaths by broad cause and leading causes 2010-2015 79% 81% 4%8% 7% <5 years 7% 9% 5% 5-14 40% 21% 17% 19% 15-24 26% 11% 15% 25-64 23% 20% 65+ 14% 3% 15% 6% 29 Broadcause Total 25% 24% 21% Injury NCD Rank Male HIV and TB Lower respiratory infections (25.5%) Lower respiratory infections (22.4%) Diarrhoeal diseases (20.1%) Diarrhoeal diseases (21.5%) Comm mat peri nut Preterm birth complications (15.7%) Preterm birth complications (17.2%) Birth asphyxia (7.0%) Birth asphyxia (8.4%) Prov, District FS, Fezile Dabi DC20 years 5 Protein-energy malnutrition (6.4%) Protein-energy malnutrition (5.7%) HIV/AIDS (4.9%) 6 HIV/AIDS (3.6%) Show history 2 Sepsis/other newborn infectious (2.2%) Sepsis/other newborn infectious (2.9%) Tuberculosis (1.9%) Road injuries (2.3%) Percentages are shown Road injuries (1.8%) Other perinatal conditions (2.0%) according to all the deaths within the age/gender Endocrine nutritional, blood, immune (1.4%) Accidental threats to breathing (1.3%) Road injuries (18.9%) Lower respiratory infections (25.3%) category of each box. Diarrhoeal diseases (17.6%) Drowning (18.3%) although only the leading Lower respiratory infections (12.2%) 10 causes are displayed. Tuberculosis (16.4%) Tuberculosis (11.7%) Road injuries (12.5%) 5 HIV/AIDS (10.6%) HIV/AIDS (9.1%) Accidental threats to breathing (6.5%) Drowning (8.2%) Endocrine nutritional, blood, immune (3.2%) Diarrhoeal diseases (4.6%) Fires, hot substances (3.1%) Fires, hot substances (4.2%) Meningitis/encephalitis (3.1%) Epilepsy (3.3%) Intestinal obstruction, strangulated he (2.7%) 10 HIV/AIDS (19.9%) Road injuries (20.5%) Tuberculosis (19.3%) Interpersonal violence (16.2%) Lower respiratory infections (14.7%) Accidental threats to breathing (10.7%) Road injuries (10.3%) Tuberculosis (10.2%) 15-24 5 Diarrhoeal diseases (5.1%) Lower respiratory infections (6.2%) 6 Interpersonal violence (4.3%) HIV/AIDS (5.7%) Endocrine nutritional, blood, immune (3.1%) Drowning (5.1%) Mechanical forces (3.8%) Meningitis/encephalitis (2.7%) Fires, hot substances (2.9%) Cardiomyopathy (2.6%) Indirect maternal (2.4%) Epilepsy (2.3%) 10 HIV/AIDS (16.6%) Tuberculosis (18.2%) Lower respiratory infections (14.0%) Tuberculosis (14.7%) Lower respiratory infections (13.7%) HIV/AIDS (12.5%) Cerebrovascular disease (5.9%) Road injuries (6.6%) 25-64 Diarrhoeal diseases (4.9%) Ischaemic heart disease (5.6%) Hypertensive heart disease (4.7%) Cerebrovascular disease (4.4%) Diabetes mellitus (4.2%) Diarrhoeal diseases (3.3%) Ischaemic heart disease (3.3%) Nephritis/nephrosis (2.6%) Endocrine nutritional, blood, immune (3.1%) Hypertensive heart disease (2.5%) 10 Road injuries (3.0%) Accidental threats to breathing (2.4%) Cerebrovascular disease (17.0%) Ischaemic heart disease (14.5%) Ischaemic heart disease (12.9%) Cerebrovascular disease (11.9%) Hypertensive heart disease (12.4%) Lower respiratory infections (10.0%) Diabetes mellitus (8.1%) Hypertensive heart disease (6.6%) Lower respiratory infections (8.0%) Prostate (5.1%) 65+ Nephritis/nephrosis (4.1%) COPD (5.1%) Diarrhoeal diseases (3.7%) Tuberculosis (5.0%) Endocrine nutritional, blood, immune (2.3%) Diabetes mellitus (5.0%) Tuberculosis (2.2%) Nephritis/nephrosis (4.1%) COPD (2.0%) Diarrhoeal diseases (2.8%) 10 Rank Maternal conditions All other causes Indirect maternal (36.3%) n=27 HIV/AIDS (21.6%) n=1 122 Other maternal (26.6%) n=19 Tuberculosis (18.5%) n=964 Hypertension in pregnancy (16.1%) n=12 Lower respiratory infections (16.1%) n=838 Maternal haemorrhage (8.8%) n=6 Diarrhoeal diseases (5.4%) n=278 Female Maternal sepsis (6.1%) n=4 Road injuries (4.5%) n=232 Abortion (6.1%) n=4 Endocrine nutritional, blood, immune (3.4%) n=175 Cerebrovascular disease (2.7%) n=143 Meningitis/encephalitis (2.7%) n=143 Nephritis/nephrosis (2.2%) n=114

Cervix (2.0%) n=102

Free State - Fezile Dabi District Municipality District Health Barometer 2016/17